let's crack defence exams www.ssbcrackexams.com

THE MOST POPULAR DEFENCE EXAM APP OF ALL TIME

Prepare for defence exams like NDA, CDS, AFCAT, INET, TA, GROUP X&Y, MNS, ACC and more.



BEST FEATURES

SSBCrackExams is the exclusive online platform for the candidates aspiring to make the Nation proud.



DETAILED LECTURES

Start enjoying your learning experience by the rock-solid video lessons specially designed for your easy understanding & higher scores in the exam.



PYQAs & TEST SERIES

Get ready for your D-Day with unlimited practice papers & previous years' questions made easy with our insightful video classes & extra notes prepared



SMART ANALYTICS

Get your own personalized Performance Tracker where you can check your Learning Footprints, Digital Notes, All-India Ranking, Timed Tests, Mind-blowing Quizzes, Prompt Videos, quick & accurate Mock Test Results, Self-Analytic Progress Reports and more.





www.ssbcrackexams.com

1. What is the modulus of the complex number 4. $\frac{\cos \theta + i \sin \theta}{\cos \theta - i \sin \theta}, \text{ where } i = \sqrt{-1}$?



(d) 2

2. Consider the proper subsets of {1, 2, 3, 4}. How many of these proper subsets are superset of the set {3} ?



3. Let p, q and r be three distinct positive real numbers. If D = $\begin{vmatrix} p & q & r \\ q & r & p \\ r & p & q \end{vmatrix}$, then which

one of the following is correct?



KJU-S-TMS

What is the sum of the last five coefficients in the expansion of $(1 + x)^9$ when it is expanded in ascending powers of x?



(c) 1024

- (d) 2048
- 5. Consider the following in respect of a non-singular matrix of order 3:
 - 1. A(adj A) = (adj A) A
 - 2. |adj A| = |A|

Which of the above statements is/are correct?

- (, 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- 6. The center of the circle (x - 2a) (x - 2b) + (y - 2c) (y - 2d) = 0 is (a) (2a, 2c) (b) (2b, 2d)
 - (c) (a + b, c + d)
 - (d) (a b, c d)
- 7. The point (1, -1) is one of the vertices of a square. If 3x + 2y = 5 is the equation of one diagonal of the square, then what is the equation of the other diagonal?

3x - 2y = 5(b) 2x - 3y = 1(c) 2x - 3y = 5(d) 2x + 3y = -1

(3-B)

- 8. Let P(x, y) be any point on the ellipse 12. $25x^2 + 16y^2 = 400$. If Q(0, 3) and R(0, -3) are two points, then what is (PQ + PR) equal to ?
 - (a) 12
 (b) 10
 (c) 8
 - (d) 6

9. If the circumcentre of the triangle formed by the lines x + 2 = 0, y + 2 = 0 and kx + y + 2 = 0 is (-1, -1), then what is the value of k?

- (a) -1
- (b) 2

1

(d) 2

10. In the parabola, $y^2 = x$, what is the length of the chord passing through the vertex and inclined to the x-axis at an angle θ ?



- (b) $\cos \theta \cdot \csc^2 \theta$
- (c) $\cot \theta \cdot \sec^2 \theta$
- (d) $2 \tan \theta \cdot \operatorname{cosec}^2 \theta$
- 11. Under which condition, are the points (a, b), (c, d) and (a - c, b - d) collinear?
 - (a) ab = cd
 - (b) ac = bd
 - c ad = bc
 - (d) abc = d



Let ABC be a triangle. If D(2, 5) and E(5, 9) are the mid-points of the sides AB and AC respectively, then what is the length of the side BC?



- 13. If the foot of the perpendicular drawn from the point (0, k) to the line 3x 4y 5 = 0 is (3, 1), then what is the value of k?
 - (a) 3
 (b) 4
 (c) 5
 (d) 6
- 14. What is the obtuse angle between the lines whose slopes are $2 \sqrt{3}$ and $2 + \sqrt{3}$?
 - (a) 105°
 - (c) 135°

 (\mathbf{b})

(5 - B)

120°

- . .
- (d) 150°
- 15. If 3x 4y 5 = 0 and 3x 4y + 15 = 0 are the equations of a pair of opposite sides of a square, then what is the area of the square?
 - (a) 4 square units
 - (b) 9 square units

) 16 square units

(d) 25 square units

Directions for the following three (03) items : Read the following information and answer the Read the following information and answer the three items that follow : Let $a \sin^2 x + b \cos^2 x = c$; $b \sin^2 y + a \cos^2 y = d$ and $p \tan x = q \tan y$.

What is $\tan^2 x$ equal to? 16.

- $\frac{c-b}{a-c}$ $\frac{a-c}{c-b}$ (b) $\frac{c-a}{c-b}$ (c)
 - (d) $\frac{c-b}{c-a}$
- 17. What is $\frac{d-a}{b-d}$ equal to?
 - (a) $\sin^2 y$ (b) $\cos^2 y$ $\tan^2 y$ (d) $\cot^2 y$
- 18. What is $\frac{p^2}{q^2}$ equal to ?
 - $\frac{(\mathbf{b}-\mathbf{c})(\mathbf{b}-\mathbf{d})}{(\mathbf{a}-\mathbf{d})(\mathbf{a}-\mathbf{c})}$ (a)
 - (b) $\frac{(a-d)(c-a)}{(b-c)(d-b)}$ (c) $\frac{(d-a)(c-a)}{(b-c)(d-b)}$

(d)
$$\frac{(b-c)(b-d)}{(c-a)(a-d)}$$

Directions for the following three (03) items :

three items that follow :

Let $t_n = \sin^n \theta + \cos^n \theta$.

What is $\frac{\mathbf{t}_3 - \mathbf{t}_5}{\mathbf{t}_5 - \mathbf{t}_7}$ equal to ? 19.



20. What is $t_1^2 - t_2$ equal to ? (a) $\cos 2\theta$ $\sin 2\theta$ (c) $2\cos\theta$ (d) $2\sin\theta$

What is the value of t_{10} where $\theta = 45^{\circ}$? 21.

1 (a) (b) $\frac{1}{32}$ (d)

(7 - B)



(9 - B)

Af $\cot \alpha$ and $\cot \beta$ are the roots of the equation Directions for the following three (03) items : $x^2 - 3x + 2 = 0$, then what is $\cot(\alpha + \beta)$ equal Consider the following Venn diagram, where X, Y and Z are three sets. Let the number of elements in to? Z be denoted by n(Z) which is equal to 90. Х Y a 16 b 18 12 17 c WB 2 (c) Z (d) 3 31) If the number of elements in Y and Z are in the ratio 4:5, then what is the value of b? The roots α and β of a quadratic equation, 29. 18 (a) satisfy the relations $\alpha + \beta = \alpha^2 + \beta^2$ and 19 (b) $\alpha\beta = \alpha^2\beta^2$. What is the number of such 21 quadratic equations? 23 (a) 0 What is the value of 2 $\mathbf{n}(\mathbf{X}) + \mathbf{n}(\mathbf{Y}) + \mathbf{n}(\mathbf{Z}) - \mathbf{n}(\mathbf{X} \cap \mathbf{Y}) - \mathbf{n}(\mathbf{Y} \cap \mathbf{Z})$ $n(X \cap Z) + n(X \cap Y \cap Z)$? (c) 3 a + b + 43(a) (d) 4 (b) a + b + 63a + b + 96(c) a + b + 106What is the argument of the complex number $\frac{1-i\sqrt{3}}{1+i\sqrt{3}}$, where $i = \sqrt{-1}$? If the number of elements belonging to neither 23. X, nor Y, nor Z is equal to p, then what is the number of elements in the complement of X ? 240° p + b + 60(b) 210° (b) p + b + 40(c) 120° (c) p + a + 60(d) 60° p + a + 40(**d**) **KJU-S-TMS** (11 - B)

Directions for the following two (02) items :	Directions for the following two (02) items :	
Read the following information and answer the two items that follow :	Read the following information and answer the two items that follow :	
Let $\frac{\tan 3A}{\tan A} = K$, where $\tan A \neq 0$ and $K \neq \frac{1}{3}$.	ABCD is a trapezium such that AB and CD are parallel and BC is perpendicular to them. Let $\angle ADB = \theta$, $\angle ABD = \alpha$, BC = p and CD = q.	
34. What is $\tan^2 A$ equal to ?	36. Consider the following : AD $\sin \theta = AB \sin \alpha$	
(a) $\frac{K+3}{3K-1}$	2.) BD sin θ = AB sin (θ + α) Which of the above is/are correct ?	
	(a) 1 only	
(b) $\frac{K-3}{2K-1}$	(b) 2 only	
3K-1	(c) Both 1 and 2	
	(d) Neither 1 nor 2	
$(c) \qquad \frac{3K-3}{K-3}$	37. What is AB equal to ?	
ASSI ASSI	(a) $\frac{(p^2 + q^2)\sin\theta}{p\cos\theta + q\sin\theta}$	
$(d) \qquad \frac{K+3}{3K+1}$	(b) $\frac{(p^2 - q^2)\cos\theta}{p\cos\theta + q\sin\theta}$	
	(c) $\frac{(p^2 + q^2)\sin\theta}{q\cos\theta + p\sin\theta}$	
35. For real values of tan A, K cannot lie between	(d) $\frac{(p^2 - q^2) \cos \theta}{q \cos \theta + p \sin \theta}$	
$\frac{1}{3}$ and 3	38. If $\tan \theta = \frac{\cos 17^\circ - \sin 17^\circ}{\cos 17^\circ + \sin 17^\circ}$, then what is the	
(b) $\frac{1}{2}$ and 2	value of θ ? (a) 0°	
(c) $\frac{1}{5}$ and 5	(b) 28° (c) 38°	
(d) $\frac{1}{7}$ and 7	(d) 52°	
KJU-S-TMS (13	– B)	

39. A and B are positive acute angles such that cos 2B = 3 sin² A and 3 sin 2A = 2 sin 2B. What is the value of (A + 2B)?



40. What is $\sin 3x + \cos 3x + 4 \sin^3 x - 3 \sin x +$

equal to ?

(c) 0
(b) 1
(c) 2 sin 2x
(d) 4 cos 4x

41. The value of ordinate of the graph of y = 2 + cos x lies in the interval

 $3\cos x - 4\cos^3 x$

- (a) [0, 1]
- (b) [0, 3]
- (c) [-1, 1]
- d [1, 3]

42. What is the value of

8 cos 10°. cos 20°. cos 40°?

- (a) tan 10°
 - cot 10°
- (c) cosec 10°
- (d) sec 10°

KJU-S-TMS

What is the value of $\cos 48^\circ - \cos 12^\circ$?



44. Consider the following statements :

- 1.) If ABC is a right-angled triangle, right-angled at A and if $\sin B = \frac{1}{3}$, then cosec C = 3.
- 2. If b cos B = c cos C and if the triangle ABC is not right-angled, then ABC must be isosceles.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- Neither 1 nor 2
- 45. Consider the following statements :
 - 1. If in a triangle ABC, A = 2B and b = c, then it must be an obtuse-angled triangle.
 - 2. There exists no triangle ABC with A = 40°, B = 65° and $\frac{a}{c} = \sin 40^{\circ} \operatorname{cosec} 15^{\circ}$.

Which of the above statements is/are correct?

(a) 1 only

b) 2 only

(c) Both 1 and 2

Neither 1 nor 2

(15 - B)



54. Let
$$A = \begin{bmatrix} x + y & y \\ 2x & x - y \end{bmatrix}$$
, $B = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$ and
 $C = \begin{bmatrix} 3 \\ 2 \end{bmatrix}$. If AB = C, then what is the value of
the determinant of the matrix A?
(a) -10
(b) -14
(c) -24
(d) -34
55. If $1.5 \le x \le 4.5$, then which one of the
following is correct?
(a) $(2x - 3)(2x - 9) \ge 0$
(b) $(2x - 3)(2x - 9) \ge 0$
(c) $(2x - 3)(2x - 9) \ge 0$
(d) $(2x - 3)(2x - 9) \ge 0$
(e) Let S = (1, 2, 3, ...). A relation R on S × S is
defined by xRy if $\log_a x \ge \log_a y$ when $a = \frac{1}{2}$.
Then the relation is
(a) reflexive only
(b) symmetric only
(c) transitive only
(d) both symmetric and transitive
57. What is the value of the determinant
 $\frac{i}{i^2} \frac{i^2}{i^3}$ where $i = \sqrt{-1}$?
(a) 0
(b) -2
(c) $4i$
(d) $4i$
(d) $-4i$
KUUS.TME (19-B)

1. 1. 1. 1.

let's crack defence exams www.ssbcrackexams.com

THE MOST POPULAR DEFENCE EXAM APP OF ALL TIME

Prepare for defence exams like NDA, CDS, AFCAT, INET, TA, GROUP X&Y, MNS, ACC and more.



BEST FEATURES

SSBCrackExams is the exclusive online platform for the candidates aspiring to make the Nation proud.



DETAILED LECTURES

Start enjoying your learning experience by the rock-solid video lessons specially designed for your easy understanding & higher scores in the exam.



PYQAs & TEST SERIES

Get ready for your D-Day with unlimited practice papers & previous years' questions made easy with our insightful video classes & extra notes prepared



SMART ANALYTICS

Get your own personalized Performance Tracker where you can check your Learning Footprints, Digital Notes, All-India Ranking, Timed Tests, Mind-blowing Quizzes, Prompt Videos, quick & accurate Mock Test Results, Self-Analytic Progress Reports and more.





www.ssbcrackexams.com

Directions for the following three (03) items :

Read the following information and answer the

Marks	Number of students	
	Physics	Mathematics
10 - 20	8	10
20 - 30	11	10
30 - 40	30	21
40 - 50	26	38
50 - 60	15	15
60 70	15	10
00 - 70	10	6

- The difference between number of students 61. under Physics and Mathematics is largest for the interval
 - 20 30(a)
 - (b) 30 - 40
 - 40 50
 - (d) **50 - 6**0

62. **Consider the following statements :**

- 1. Modal value of the marks in Physics lies in the interval 30 - 40.
- Median of the marks in Physics is less 2. than that of marks in Mathematics.

Which of the above statements is/are correct?

(a) 1 only

- (b) 2 only
- Both 1 and 2 (c)
- Neither 1 nor 2 (d)

What is the mean of marks in Physics ? 63.



KJU-S-TMS

What is the standard deviation of the 64. observations



- If $\sum x_i = 20$, $\sum x_i^2 = 200$ and n = 10 for an 65. observed variable x, then what is the coefficient of variation ?
 - 80 (a)



What is the probability that February of a 66. leap year selected at random, will have five Sundays?

(d)

(a)

(c)

1

(21 - B)

- 67. The arithmetic mean of 100 observations is 40. Later, it was found that an observation '53' was wrongly read as '83'. What is the correct arithmetic mean ?
 - (a) 39·8
 - (b) 39·7
 - (c) 39·6
 - (d) 39.5
- **68.** A husband and wife appear in an intervew for two vacancies for the same post. The probability of the husband's selection is $\frac{1}{7}$ and that of the wife's selection is $\frac{1}{5}$. If the events are independent, then the probability of which one of the following is $\frac{11}{35}$?

(a) At least one of them will be selected

- (b) Only one of them will be selected
- (c) None of them will be selected
- (d) Both of them will be selected
- 69. A dealer has a stock of 15 gold coins out of which 6 are counterfeits. A person randomly picks 4 of the 15 gold coins. What is the probability that all the coins picked will be counterfeits?



70. A committee of 3 is to be formed from a group of 2 boys and 2 girls. What is the probability that the committee consists of 2 boys and 1 girl?



Crack EXAMS

71. In a lottery of 10 tickets numbered 1 to 10, two tickets are drawn simultaneously. What is the probability that both the tickets drawn have prime numbers ?



KJU-S-TMS

(23 – B)

- **72.** Let X and Y represent prices (in \mathfrak{P}) of a commodity in Kolkata and Mumbai respectively. It is given that $\overline{X} = 65$, $\overline{Y} = 67$, $\sigma_X = 2.5$, $\sigma_Y = 3.5$ and r(X, Y) = 0.8. What is the equation of regression of Y on X?
 - (a) Y = 0.175X 5
 - (b) Y = 1.12X 5.8
 - (c) Y = 1.12X 5
 - (d) Y = 0.17X + 5.8
- 73. Consider a random variable X which follows Binomial distribution with parameters n = 10and $p = \frac{1}{5}$. Then Y = 10 - X follows Binomial distribution with parameters n and p respectively given by
 - (a) 5, $\frac{1}{5}$
 - (b) 5, $\frac{2}{5}$
 - (c) 10, $\frac{3}{5}$

If A and B are two events such that P(A) = 0.6, P(B) = 0.5 and $P(A \cap B) = 0.4$, then consider the following statements : P(B) = 0.6

- 1. $P(\overline{A} \cup B) = 0.9$.
- 2. $P(\overline{B} | \overline{A}) = 0.6$.

Which of the above statements is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- 75. Three cooks X, Y and Z bake a special kind of cake, and with respective probabilities 0.02, 0.03 and 0.05, it fails to rise. In the restaurant where they work, X bakes 50%, Y bakes 30% and Z bakes 20% of cakes. What is the proportion of failures caused by X ?



(25 – B)



KJU-S-TMS

(27 – B)

- 82. Consider the following statements :
 - 1. The function f(x) = ln x increases in the interval $(0, \infty)$.
 - 2. The function $f(x) = \tan x$ increases in the interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

Which of the above statements is/are correct?

- (a) 1 only
 - 🖉 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2
- 83. Which one of the following is correct in respect of the graph of $y = \frac{1}{x-1}$?
 - (a) The domain is $\{x \in \mathbb{R} | x \neq 1\}$ and the range is the set of reals.

The domain is $\{x \in \mathbf{R} \mid x \neq 1\}$, the range is $\{y \in \mathbf{R} \mid y \neq 0\}$ and the graph intersects y-axis at (0, -1).

- (c) The domain is the set of reals and the range is the singleton set {0}.
- (d) The domain is $\{x \in \mathbf{R} | x \neq 1\}$ and the range is the set of points on the y-axis.
- 84. What is the solution of the differential equation $ln\left(\frac{dy}{dx}\right) = x$?

(a) $y = e^{x} + c$ (b) $y = e^{-x} + c$ (c) y = ln x + c(d) y = 2 ln x + c 85. Let l be the length and b be the breadth of a rectangle such that l + b = k. What is the maximum area of the rectangle ?



86. The numbers 4 and 9 have frequencies x and (x - 1) respectively. If their arithmetic mean is 6, then what is the value of x ?



If three dice are rolled under the condition that no two dice show the same face, then what is the probability that one of the faces is having the number 6 ?



(29 – B)

88. If $P(A \cup B) = \frac{5}{6}$, $P(A \cap B) = \frac{1}{3}$ and $P(\text{not } A) = \frac{1}{2}$, then which one of the following is **not** correct ? (a) $P(B) = \frac{2}{3}$ (b) $P(A \cap B) = P(A)P(B)$ (c) $P(A \cup B) > P(A) + P(B)$ (d) P(not A and not B) = P(not A) P(not B)

- 89. The sum of deviations of n number of observations measured from 2.5 is 50. The sum of deviations of the same set of observations measured from 3.5 is 50. What is the value of n?
 - (a) 50
 - (b) 60
 - (c) 80
 - (d) 100
- 90. A data set of n observations has mean 2M, while another data set of 2n observations has mean M. What is the mean of the combined data sets ?
 - (a) M
 - (b) $\frac{3M}{2}$
 - (c) $\frac{2M}{3}$
 - $(d) \quad \frac{4M}{3}$

91. If $f(x) = 3x^2 - 5x + p$ and f(0) and f(1) are opposite in sign, then which of the following is correct?



- 92. If $e^{\theta \phi} = c + 4\theta \phi$, where c is an arbitrary constant and ϕ is a function of θ , then what is $\phi d\theta$ equal to ?
 - (a) $\theta d\phi$

(b)
$$-\theta d\phi$$

(c) $4\theta d\phi$
(d) $-4\theta d\phi$

93. If $p(x) = (4e)^{2x}$, then what is $\int p(x) dx$ equal to?

(a) $\frac{p(x)}{1+2 \ln 2} + c$

(b)
$$\frac{p(x)}{2(1+2\ln 2)} + c$$

(c)
$$\frac{2p(x)}{1+ln 4} + c$$

$$\frac{\mathbf{p}(\mathbf{x})}{1+\ln 2} + \mathbf{c}$$

KJU-S-TMS

(31 – B)

94. What is the value of
$$\int_{0}^{\pi/4} (\tan^{3} x + \tan x) dx$$
?
(a) $\frac{1}{4}$
(b) $\frac{1}{2}$
(c) 1
(d) 2
95. Let $y = 3x^{2} + 2$. If x changes from 10 to 10-1, then what is the total change in y?
(a) 4.71
(b) 5.23
(c) 6.03
(c) 6.01
96. If $f(x) = \frac{\sin x}{x}$, where $x \in \mathbf{B}$, is to be continuous at $x = 0$, then the value of the function $f(x) = \cos^{-1}(x-2)$?
(a) $should be 0$
(b) $should be 1$
(c) $should be 2$
(d) cannot be determined
(c) $(1 + y^{2}) dx$ is
(c) $1 + y^{2} dx$ is
(a) $y = \tan x + c$
(b) $y = \tan (x + c)$
(c) $\tan^{-1}(y + c) = x$
(d) $\tan^{-1}(y + c) = 2x$
(e) $\tan^{-1}(y + c) = 2x$
(f) $\tan^{-1}(y + c) = 2x$
(g) $\tan x + x \cos x + \frac{\sin^{2} x}{2} + c$
(h) $\sin x - x \cos x + \frac{\sin^{2} x}{2} + c$
(h) $x \sin x - x \cos x + \frac{\sin^{2} x}{2} + c$
(h) $x \sin x - x \cos x + \frac{\sin^{2} x}{2} + c$
(h) $x \sin x - x \cos x + \frac{\sin^{2} x}{2} + c$
(h) $(1, 3)$
(h) $(1, 2, 1]$
(h) $(1, 2, 1]$
(h) $(1, 2, 1]$

KJU-S-TMS

(33 – ^B)

100. What is the area of the region enclosed between the curve $y^2 = 2x$ and the straight 103. What is $\int \frac{dx}{x(x^n+1)}$ equal to ? line y = x? (a) $\frac{1}{n} ln \left(\frac{\mathbf{x}^n}{\mathbf{x}^n + 1} \right) + c$ 1 2 (a) (b) $ln\left(\frac{\mathbf{x}^n+1}{\mathbf{x}^n}\right)+\mathbf{c}$ (b) 1 23 (c) $ln\left(\frac{x^n}{x^n+1}\right)+c$ (d) 2 (d) $\frac{1}{n}ln\left(\frac{x^{n}+1}{x^{n}}\right)+c$ 101. If $f(x) = 2x - x^2$, then what is the value of f(x + 2) + f(x - 2) when x = 0? 104. What is the minimum value of |x - 1|, where - 8 $\mathbf{x} \in \mathbf{R}$? (b) - 4 (a) 0 (c) 8 (b) 1 (c) (d) 4 102. If $x^m y^n = a^{m+n}$, then what is $\frac{dy}{dx}$ equal to ? 105. What is the value of k such that integration of $\frac{3x^2+8-4k}{x}$ with respect to x, may be a my nx (a) rational function ? my (a) 0 nx (b) mx (c) nv $\frac{ny}{mx}$ (**d**) (**d**) -2

KJU-S-TMS

(35 – B)

106. What is the length of the diameter of the 110. What is the equation of the plane which cuts sphere whose centre is at (1, -2, 3) and which touches the plane 6x - 3y + 2z - 4 = 0?

- (a) 1 unit
- (b) 2 units
- (c) 3 units



- 107. What is the perpendicular distance from the point (2, 3, 4) to the line $\frac{x-0}{1} = \frac{y-0}{0} = \frac{z-0}{0}$?
 - (a) 6 units

5 units

- 3 units (c)
- (d) 2 units
- 108. If a line has direction ratios < a + b, b + c, c + a >, then what is the sum of the squares of its direction cosines?
 - $(a + b + c)^2$ (a)
 - 2(a + b + c)(b)
 - 3 (c)

1

- (\mathbf{d})
- many compartments do the 109. Into how coordinate planes divide the space ?
 - 2 (a) (b) 4 8 (d) 16
- KJU-S-TMS

an intercept 5 units on the z-axis and is parallel to xy-plane?

(a)
$$x + y = 5$$

(b) $z = 5$
(c) $z = 0$
(d) $x + y + z = 5$

111. If a is a unit vector in the xy-plane making an angle 30° with the positive x-axis, then what is \hat{a} equal to?

 $\frac{\sqrt{3}i+j}{2}$ (b) $\frac{\sqrt{3}\hat{i}-\hat{j}}{2}$ $\frac{\hat{i}+\sqrt{3}\hat{j}}{2}$ (c) $\frac{\hat{i}-\sqrt{3}\hat{j}}{2}$ (d)

112. Let A be a point in space such that $|\overrightarrow{OA}| = 12$, where O is the origin. If \overrightarrow{OA} is inclined at angles 45° and 60° with x-axis and y-axis respectively, then what is \overrightarrow{OA} equal to?

(a)
$$6\hat{i} + 6\hat{j} \pm \sqrt{2}\hat{k}$$

(b) $6\hat{i} + 6\sqrt{2}\hat{j} \pm 6\hat{k}$
(c) $6\sqrt{2}\hat{i} + 6\hat{j} \pm 6\hat{k}$
(d) $3\sqrt{2}\hat{i} + 3\hat{j} \pm 6\hat{k}$

(37 - B)



KJU-S-TMS

(39 - B)

let's crack defence exams www.ssbcrackexams.com

THE MOST POPULAR DEFENCE EXAM APP OF ALL TIME

Prepare for defence exams like NDA, CDS, AFCAT, INET, TA, GROUP X&Y, MNS, ACC and more.



BEST FEATURES

SSBCrackExams is the exclusive online platform for the candidates aspiring to make the Nation proud.



DETAILED LECTURES

Start enjoying your learning experience by the rock-solid video lessons specially designed for your easy understanding & higher scores in the exam.



PYQAs & TEST SERIES

Get ready for your D-Day with unlimited practice papers & previous years' questions made easy with our insightful video classes & extra notes prepared



SMART ANALYTICS

Get your own personalized Performance Tracker where you can check your Learning Footprints, Digital Notes, All-India Ranking, Timed Tests, Mind-blowing Quizzes, Prompt Videos, quick & accurate Mock Test Results, Self-Analytic Progress Reports and more.





👼 www.ssbcrackexams.com