

1. Consider the following statements :

1.  $f(x) = \ln x$  is increasing in  $(0, \infty)$

2.  $g(x) = e^x + e^{\frac{1}{x}}$  is decreasing in  $(0, \infty)$

Which of the statements given above is/are correct ?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2

2. What is the derivative of  $\sin^2 x$  with respect to  $\cos^2 x$  ?

- (a) -1
- (b) 1
- (c)  $\sin 2x$
- (d)  $\cos 2x$

3. For what value of  $m$  with  $m < 0$ , is the area bounded by the lines  $y = x$ ,  $y = mx$  and  $x = 2$  equal to 3 ?

- (a)  $-\frac{1}{2}$
- (b) -1
- (c)  $-\frac{3}{2}$
- (d) -2

4. What is the derivative of  $\operatorname{cosec}(x^\circ)$  ?

- (a)  $-\operatorname{cosec}(x^\circ) \cot(x^\circ)$
- (b)  $-\frac{\pi}{180} \operatorname{cosec}(x^\circ) \cot(x^\circ)$

(c)  $\frac{\pi}{180} \operatorname{cosec}(x^\circ) \cot(x^\circ)$

(d)  $-\frac{\pi}{180} \operatorname{cosec}(x) \cot(x)$

5. A solution of the differential equation

$$\left(\frac{dy}{dx}\right)^2 - x \frac{dy}{dx} = 0 \text{ is}$$

- (a)  $y = 2x$
- (b)  $y = 2x + 4$
- (c)  $y = x^2 - 1$
- (d)  $y = \frac{(x^2 - 2)}{2}$

6. If  $f(x) = x^2 + 2$  and  $g(x) = 2x - 3$ , then what is  $(fg)(1)$  equal to ?

- (a) 3
- (b) 1
- (c) -2
- (d) -3

7. What is the range of the function  $f(x) = x + |x|$  if the domain is the set of real numbers ?

- (a)  $(0, \infty)$
- (b)  $[0, \infty)$
- (c)  $(-\infty, \infty)$
- (d)  $[1, \infty)$

8. If  $f(x) = x(4x^2 - 3)$ , then what is  $f(\sin\theta)$  equal to?

- (a)  $-\sin 3\theta$
- (b)  $-\cos 3\theta$
- (c)  $\sin 3\theta$
- (d)  $-\sin 4\theta$

9. What is  $\lim_{x \rightarrow 5} \frac{5-x}{|x-5|}$  equal to?

- (a)  $-1$
- (b)  $0$
- (c)  $1$
- (d) Limit does not exist

10. What is  $\lim_{x \rightarrow 1} \frac{x^9 - 1}{x^3 - 1}$  equal to?

- (a)  $-1$
- (b)  $-3$
- (c)  $3$
- (d) Limit does not exist

Consider the following for the next three (03) items that follow:

Let  $f(x) = Pe^x + Qe^{2x} + Re^{3x}$ , where  $P, Q, R$  are real numbers. Further  $f(0) = 6$ ,  $f'(\ln 3) = 282$  and  $\int_0^{\ln 2} f(x) dx = 11$

11. What is the value of  $Q$ ?

- (a)  $1$
- (b)  $2$

(c)  $3$

(d)  $4$

12. What is the value of  $R$ ?

- (a)  $1$
- (b)  $2$
- (c)  $3$
- (d)  $4$

13. What is  $f'(0)$  equal to?

- (a)  $18$
- (b)  $16$
- (c)  $15$
- (d)  $14$

Consider the following for the next two (02) items that follow:

Suppose  $E$  is the differential equation representing family of curves  $y^2 = 2cx + 2c\sqrt{c}$  where  $c$  is a positive parameter.

14. What is the order of the differential equation?

- (a)  $1$
- (b)  $2$
- (c)  $3$
- (d)  $4$

15. What is the degree of the differential equation ?

- (a) 2
- (b) 3
- (c) 4
- (d) Degree does not exist

Consider the following for the next three (03) items that follow :

$$\text{Let } f(x) = \begin{vmatrix} \cos x & x & 1 \\ 2 \sin x & x^2 & 2x \\ \tan x & x & 1 \end{vmatrix}$$

16. What is  $f(0)$  equal to ?

- (a) -1
- (b) 0
- (c) 1
- (d) 2

17. What is  $\lim_{x \rightarrow 0} \frac{f(x)}{x}$  equal to ?

- (a) -1
- (b) 0
- (c) 1
- (d) 2

18. What is  $\lim_{x \rightarrow 0} \frac{f(x)}{x^2}$  equal to ?

- (a) -1
- (b) 0
- (c) 1
- (d) 2

Consider the following for the next two (02) items that follow :

Let  $f(x) = \sin[\pi^2]x + \cos[-\pi^2]x$  where  $[\cdot]$  is a greatest integer function

19. What is  $f\left(\frac{\pi}{2}\right)$  equal to ?

- (a) -1
- (b) 0
- (c) 1
- (d) 2

20. What is  $f\left(\frac{\pi}{4}\right)$  equal to ?

- (a)  $-\frac{1}{\sqrt{2}}$
- (b) -1
- (c) 1
- (d)  $\frac{1}{\sqrt{2}}$

Consider the following for the next two (02) items that follow :

$$\text{Let } \Delta(a, b, c, \alpha) = \begin{vmatrix} a & b & a\alpha + b \\ b & c & b\alpha + c \\ a\alpha + b & b\alpha + c & 0 \end{vmatrix}$$

21. If  $\Delta(a, b, c, \alpha) = 0$  for every  $\alpha > 0$ , then which one of the following is correct ?

- (a)  $a, b, c$  are in AP
- (b)  $a, b, c$  are in GP
- (c)  $a, 2b, c$  are in AP
- (d)  $a, 2b, c$  are in GP

22. If  $\Delta(7, 4, 2, \alpha) = 0$ , then  $\alpha$  is a root of which one of the following equations ?

- (a)  $7x^2 + 4x + 2 = 0$
- (b)  $7x^2 - 4x + 2 = 0$
- (c)  $7x^2 + 8x + 2 = 0$
- (d)  $7x^2 - 8x + 2 = 0$

Consider the following for the next two (02) items that follow :

Given that  $m(\theta) = \cot^2\theta + n^2 \tan^2\theta + 2n$ , where  $n$  is a fixed positive real number.

23. What is the least value of  $m(\theta)$  ?

- (a)  $n$
- (b)  $2n$

(c)  $3n$

(d)  $4n$

24. Under what condition does  $m$  attain the least value ?

- (a)  $n = \tan^2\theta$
- (b)  $n = \cot^2\theta$
- (c)  $n = \sin^2\theta$
- (d)  $n = \cos^2\theta$

Consider the following for the next two (02) items that follow :

A quadrilateral is formed by the lines  $x = 0, y = 0, x + y = 1$  and  $6x + y = 3$ .

25. What is the equation of diagonal through origin ?

- (a)  $3x + y = 0$
- (b)  $2x + 3y = 0$
- (c)  $3x - 2y = 0$
- (d)  $3x + 2y = 0$

26. What is the equation of other diagonal ?

- (a)  $x + 2y - 1 = 0$
- (b)  $x - 2y - 1 = 0$
- (c)  $2x + y + 1 = 0$
- (d)  $2x + y - 1 = 0$

Consider the following for the next two (02) items that follow :

$P(x, y)$  is any point on the ellipse  $x^2 + 4y^2 = 1$ . Let  $E, F$  be the foci of the ellipse.

27. What is  $PE + PF$  equal to ?

- (a) 1
- (b) 2
- (c) 3
- (d) 4

28. Consider the following points :

1.  $\left(\frac{\sqrt{3}}{2}, 0\right)$

2.  $\left(\frac{\sqrt{3}}{2}, \frac{1}{4}\right)$

3.  $\left(\frac{\sqrt{3}}{2}, -\frac{1}{4}\right)$

Which of the above points lie on latus rectum of ellipse ?

- (a) 1 and 2 only
- (b) 2 and 3 only
- (c) 1 and 3 only
- (d) 1, 2 and 3

Consider the following for the next two (02) items that follow :

The line  $y = x$  partitions the circle

$$(x - a)^2 + y^2 = a^2 \text{ in two segments.}$$

29. What is the area of minor segment ?

(a)  $\frac{(\pi - 2)a^2}{4}$

(b)  $\frac{(\pi - 1)a^2}{4}$

(c)  $\frac{(\pi - 2)a^2}{2}$

(d)  $\frac{(\pi - 1)a^2}{2}$

30. What is the area of major segment ?

(a)  $\frac{(3\pi - 2)a^2}{4}$

(b)  $\frac{(3\pi + 2)a^2}{4}$

(c)  $\frac{(3\pi - 2)a^2}{2}$

(d)  $\frac{(3\pi + 2)a^2}{2}$

31. Consider the following frequency distribution :

$x$	1	2	3	5
$f$	4	6	9	7

What is the value of median of the distribution ?

- (a) 1
- (b) 2
- (c) 3
- (d) 3.5

32. For data  $-1, 1, 4, 3, 8, 12, 17, 19, 9, 11$ ; if  $M$  is the median of first 5 observations and  $N$  is the median of last five observations, then what is the value of  $4M - N$ ?

- (a) 7
- (b) 4
- (c) 1
- (d) 0

33. Let  $P, Q, R$  represent mean, median and mode. If for some distribution  $5P = 4Q = \frac{R}{2}$ , then what is  $\frac{P+Q}{2P+0.7R}$  equal to?

- (a)  $\frac{1}{12}$
- (b)  $\frac{1}{7}$
- (c)  $\frac{2}{9}$
- (d)  $\frac{1}{4}$

34. If  $G$  is the geometric mean of numbers  $1, 2, 2^2, 2^3, \dots, 2^{n-1}$ , then what is the value of  $1 + 2\log_2 G$ ?

- (a) 1
- (b) 4
- (c)  $n - 1$
- (d)  $n$

35. If  $H$  is the harmonic mean of numbers  $1, 2, 2^2, 2^3, \dots, 2^{n-1}$ , then what is  $n/H$  equal to?

- (a)  $2 - \frac{1}{2^{n+1}}$
- (b)  $2 - \frac{1}{2^{n-1}}$
- (c)  $2 + \frac{1}{2^{n-1}}$
- (d)  $2 - \frac{1}{2^n}$

36. Let  $P$  be the median,  $Q$  be the mean and  $R$  be the mode of observations  $x_1, x_2, x_3, \dots, x_n$ . Let  $S = \sum_{i=1}^n (2x_i - a)^2$ .  $S$  takes minimum value, when  $a$  is equal to

- (a)  $P$
- (b)  $\frac{Q}{2}$
- (c)  $2Q$
- (d)  $R$

37. One bag contains 3 white and 2 black balls, another bag contains 2 white and 3 black balls. Two balls are drawn from the first bag and put it into the second bag and then a ball is drawn from the second bag. What is the probability that it is white?

- (a)  $\frac{6}{7}$
- (b)  $\frac{33}{70}$
- (c)  $\frac{3}{10}$
- (d)  $\frac{1}{70}$

38. Three dice are thrown. What is the probability that each face shows only multiples of 3 ?

(a)  $\frac{1}{9}$

(b)  $\frac{1}{18}$

(c)  $\frac{1}{27}$

(d)  $\frac{1}{3}$

39. What is the probability that the month of December has 5 Sundays ?

(a) 1

(b)  $\frac{1}{4}$

(c)  $\frac{3}{7}$

(d)  $\frac{2}{7}$

40. A natural number  $n$  is chosen from the first 50 natural numbers. What is the probability that  $n + \frac{50}{n} < 50$  ?

(a)  $\frac{23}{25}$

(b)  $\frac{47}{50}$

(c)  $\frac{24}{25}$

(d)  $\frac{49}{50}$

41. How many real numbers satisfy the equation  $|x-4| + |x-7| = 15$  ?

(a) Only one

(b) Only two

(c) Only three

(d) Infinitely many

42. A mapping  $f: A \rightarrow B$  defined as  $f(x) = \frac{2x+3}{3x+5}$ ,  $x \in A$ . If  $f$  is to be onto, then what are  $A$  and  $B$  equal to ?

(a)  $A = R \setminus \{-\frac{5}{3}\}$  and  $B = R \setminus \{-\frac{2}{3}\}$

(b)  $A = R$  and  $B = R \setminus \{-\frac{5}{3}\}$

(c)  $A = R \setminus \{-\frac{3}{2}\}$  and  $B = R \setminus \{0\}$

(d)  $A = R \setminus \{-\frac{5}{3}\}$  and  $B = R \setminus \{\frac{2}{3}\}$

43.  $\alpha$  and  $\beta$  are distinct real roots of the quadratic equation  $x^2 + ax + b = 0$ . Which of the following statements is/are sufficient to find  $\alpha$  ?

1.  $\alpha + \beta = 0$ ,  $\alpha^2 + \beta^2 = 2$

2.  $\alpha\beta^2 = -1$ ,  $a = 0$

Select the correct answer using the code given below :

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

44. If the sixth term in the binomial expansion of  $\left(x^{-\frac{8}{3}} + x^2 \log_{10} x\right)^8$  is 5600,

then what is the value of  $x$ ?

- (a) 6
- (b) 8
- (c) 9
- (d) 10

45. How many terms are there in the expansion of  $(3x - y)^4(x + 3y)^4$ ?

- (a) 9
- (b) 12
- (c) 15
- (d) 17

46.  $p, q, r$  and  $s$  are in AP such that  $p + s = 8$  and  $qr = 15$ . What is the difference between largest and smallest numbers?

- (a) 6
- (b) 5
- (c) 4
- (d) 3

47. Consider the following statements for a fixed natural number  $n$ :

1.  $C(n, r)$  is greatest if  $n = 2r$
2.  $C(n, r)$  is greatest if  $n = 2r - 1$  and  $n = 2r + 1$

Which of the statements given above is/are correct?

- (a) 1 only
- (b) 2 only
- (c) Both 1 and 2
- (d) Neither 1 nor 2

48.  $m$  parallel lines cut  $n$  parallel lines giving rise to 60 parallelograms. What is the value of  $(m + n)$ ?

- (a) 6
- (b) 7
- (c) 8
- (d) 9

49. Let  $x$  be the number of permutations of the word 'PERMUTATIONS' and  $y$  be the number of permutations of the word 'COMBINATIONS'. Which one of the following is correct?

- (a)  $x = y$
- (b)  $y = 2x$
- (c)  $x = 4y$
- (d)  $y = 4x$

50. 5-digit numbers are formed using the digits 0, 1, 2, 4, 5 without repetition. What is the percentage of numbers which are greater than 50,000?

- (a) 20%
- (b) 25%
- (c)  $\frac{100}{3}\%$
- (d)  $\frac{110}{3}\%$



51. If  $2 - i\sqrt{3}$  where  $i = \sqrt{-1}$  is a root of the equation  $x^2 + ax + b = 0$ , then what is the value of  $(a + b)$ ?

- (a) -11
- (b) -3
- (c) 0
- (d) 3

52. If  $z = \frac{1+i\sqrt{3}}{1-i\sqrt{3}}$  where  $i = \sqrt{-1}$ , then what is the argument of  $z$ ?

- (a)  $\frac{\pi}{3}$
- (b)  $\frac{2\pi}{3}$
- (c)  $\frac{4\pi}{3}$
- (d)  $\frac{5\pi}{6}$

53. If  $a, b, c$  are in AP, then what is

$$\begin{vmatrix} x+1 & x+2 & x+3 \\ x+2 & x+3 & x+4 \\ x+a & x+b & x+3 \end{vmatrix} \text{ equal to ?}$$

- (a) -1
- (b) 0
- (c) 1
- (d) 2

54. If  $\log_x a$ ,  $a^x$  and  $\log_b x$  are in GP, then what is  $x$  equal to?

- (a)  $\log_a(\log_b a)$
- (b)  $\log_b(\log_a b)$
- (c)  $\frac{\log_a(\log_b a)}{2}$
- (d)  $\frac{\log_b(\log_a b)}{2}$

55. If  $2^{\frac{1}{c}}$ ,  $2^{\frac{b}{ac}}$ ,  $2^{\frac{1}{a}}$  are in GP, then which one of the following is correct?

- (a)  $a, b, c$  are in AP
- (b)  $a, b, c$  are in GP
- (c)  $a, b, c$  are in HP
- (d)  $ab, bc, ca$  are in AP

56. The first and the second terms of an AP are  $\frac{5}{2}$  and  $\frac{23}{12}$  respectively. If  $n^{\text{th}}$  term is the largest negative term, what is the value of  $n$ ?

- (a) 5
- (b) 6
- (c) 7
- (d)  $n$  cannot be determined

57. For how many integral values of  $k$ , the equation  $x^2 - 4x + k = 0$ , where  $k$  is an integer has real roots and both of them lie in the interval  $(0, 5)$ ?

- (a) 3
- (b) 4
- (c) 5
- (d) 6

58. In an AP, the first term is  $x$  and the sum of the first  $n$  terms is zero. What is the sum of next  $m$  terms?

(a)  $\frac{mx(m+n)}{n-1}$

(b)  $\frac{mx(m+n)}{1-n}$

(c)  $\frac{nx(m+n)}{m-1}$

(d)  $\frac{nx(m+n)}{1-m}$

59. Consider the following statements:

1.  $(25)! + 1$  is divisible by 26

2.  $(6)! + 1$  is divisible by 7

Which of the above statements is/are correct?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

60. If  $z$  is a complex number such that

$\frac{z-1}{z+1}$  is purely imaginary, then what is

$|z|$  equal to?

(a)  $\frac{1}{2}$

(b)  $\frac{2}{3}$

(c) 1

(d) 2

61. If  $\omega$  is a non-real cube root of 1, then

what is the value of  $\left| \frac{1-\omega}{\omega+\omega^2} \right|$ ?

(a)  $\sqrt{3}$

(b)  $\sqrt{2}$

(c) 1

(d)  $\frac{4}{\sqrt{3}}$

62. What is the number of 6-digit numbers that can be formed only by using 0, 1, 2, 3, 4 and 5 (each once); and divisible by 6?

(a) 96

(b) 120

(c) 192

(d) 312

63. What is the binary number equivalent to decimal number 1011?

(a) 1011

(b) 111011

(c) 11111001

(d) 111110011

64. Let  $A$  be a matrix of order  $3 \times 3$  and  $|A| = 4$ . If  $|2 \operatorname{adj}(3A)| = 2^\alpha 3^\beta$ , then what is the value of  $(\alpha + \beta)$ ?

(a) 12

(b) 13

(c) 17

(d) 24

65. If  $\alpha$  and  $\beta$  are the distinct roots of equation  $x^2 - x + 1 = 0$ , then what is the value of  $\left| \frac{\alpha^{100} + \beta^{100}}{\alpha^{100} - \beta^{100}} \right|$  ?

(a)  $\sqrt{3}$

(b)  $\sqrt{2}$

(c) 1

(d)  $\frac{1}{\sqrt{3}}$

66. Let  $A$  and  $B$  be symmetric matrices of same order, then which one of the following is correct regarding  $(AB - BA)$  ?

1. Its diagonal entries are equal but nonzero

2. The sum of its non-diagonal entries is zero

Select the correct answer using the code given below :

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

67. Consider the following statements in respect of square matrices  $A, B, C$  each of same order  $n$  :

1.  $AB = AC \Rightarrow B = C$  if  $A$  is non-singular

2. If  $BX = CX$  for every column matrix  $X$  having  $n$  rows then  $B = C$

Which of the statements given above is/are correct ?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

68. The system of linear equations  $x + 2y + z = 4$ ,  $2x + 4y + 2z = 8$  and  $3x + 6y + 3z = 10$  has

(a) a unique solution

(b) infinite many solutions

(c) no solution

(d) exactly three solutions

69. Let  $AX = B$  be a system of 3 linear equations with 3-unknowns. Let  $X_1$  and  $X_2$  be its two distinct solutions. If the combination  $aX_1 + bX_2$  is a solution of  $AX = B$ ; where  $a, b$  are real numbers, then which one of the following is correct ?

(a)  $a = b$

(b)  $a + b = 1$

(c)  $a + b = 0$

(d)  $a - b = 1$

70. What is the sum of the roots of the

$$\text{equation } \begin{vmatrix} 0 & x-a & x-b \\ 0 & 0 & x-c \\ x+b & x+c & 1 \end{vmatrix} = 0 ?$$

- (a)  $a + b + c$
- (b)  $a - b + c$
- (c)  $a + b - c$
- (d)  $a - b - c$

Consider the following for the next two (02) items that follow :

Let  $A(1, -1, 2)$  and  $B(2, 1, -1)$  be the end points of the diameter of the sphere  $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz - 1 = 0$ .

71. What is  $u + v + w$  equal to ?

- (a)  $-2$
- (b)  $-1$
- (c)  $1$
- (d)  $2$

72. If  $P(x, y, z)$  is any point on the sphere, then what is  $PA^2 + PB^2$  equal to ?

- (a)  $15$
- (b)  $14$
- (c)  $13$
- (d)  $6.5$

Consider the following for the next two (02) items that follow :

Consider two lines whose direction ratios are  $(2, -1, 2)$  and  $(k, 3, 5)$ . They are inclined at an angle  $\frac{\pi}{4}$ .

73. What is the value of  $k$  ?

- (a)  $4$
- (b)  $2$
- (c)  $1$
- (d)  $-1$

74. What are the direction ratios of a line which is perpendicular to both the lines ?

- (a)  $(1, 2, 10)$
- (b)  $(-1, -2, 10)$
- (c)  $(11, 12, -10)$
- (d)  $(11, 2, -10)$

Consider the following for the next two (02) items that follow :

Let  $\vec{a} = 3\hat{i} + 3\hat{j} + 3\hat{k}$  and  $\vec{c} = \hat{j} - \hat{k}$ . Let  $\vec{b}$  be such that  $\vec{a} \cdot \vec{b} = 27$  and  $\vec{a} \times \vec{b} = 9\vec{c}$

75. What is  $\vec{b}$  equal to ?

- (a)  $3\hat{i} + 4\hat{j} + 2\hat{k}$
- (b)  $5\hat{i} + 2\hat{j} + 2\hat{k}$
- (c)  $5\hat{i} - 2\hat{j} + 6\hat{k}$
- (d)  $3\hat{i} + 3\hat{j} + 4\hat{k}$

76. What is the angle between  $(\vec{a} + \vec{b})$  and  $\vec{c}$  ?

(a)  $\frac{\pi}{2}$

(b)  $\frac{\pi}{3}$

(c)  $\frac{\pi}{4}$

(d)  $\frac{\pi}{6}$

Consider the following for the next two (02) items that follow :

Let a vector  $\vec{a} = 4\hat{i} - 8\hat{j} + \hat{k}$  make angles  $\alpha, \beta, \gamma$  with the positive directions of  $x, y, z$  axes respectively.

77. What is  $\cos\alpha$  equal to ?

(a)  $\frac{1}{3}$

(b)  $\frac{4}{9}$

(c)  $\frac{5}{9}$

(d)  $\frac{2}{3}$

78. What is  $\cos 2\beta + \cos 2\gamma$  equal to ?

(a)  $-\frac{32}{81}$

(b)  $-\frac{16}{81}$

(c)  $\frac{16}{81}$

(d)  $\frac{32}{81}$

Consider the following for the next two (02) items that follow :

The position vectors of two points  $A$  and  $B$  are  $\hat{i} - \hat{j}$  and  $\hat{j} + \hat{k}$  respectively.

79. Consider the following points :

1.  $(-1, -3, 1)$

2.  $(-1, 3, 2)$

3.  $(-2, 5, 3)$

Which of the above points lie on the line joining  $A$  and  $B$  ?

(a) 1 and 2 only

(b) 2 and 3 only

(c) 1 and 3 only

(d) 1, 2 and 3

80. What is the magnitude of  $\overline{AB}$  ?

(a) 2

(b) 3

(c)  $\sqrt{6}$

(d)  $\sqrt{3}$

81. The mean and variance of five observations are 14 and 13.2 respectively. Three of the five observations are 11, 16 and 20. What are the other two observations?

- (a) 8 and 15
- (b) 9 and 14
- (c) 10 and 13
- (d) 11 and 12

82. Let  $A$  and  $B$  be two independent events such that

$$P(\bar{A}) = 0.7, P(\bar{B}) = k, P(A \cup B) = 0.8.$$

What is the value of  $k$ ?

- (a)  $\frac{5}{7}$
- (b)  $\frac{4}{7}$
- (c)  $\frac{2}{7}$
- (d)  $\frac{1}{7}$

83. A biased coin with the probability of getting head equal to  $\frac{1}{4}$  is tossed five times. What is the probability of getting tail in all the first four tosses followed by head?

- (a)  $\frac{81}{512}$
- (b)  $\frac{81}{1024}$

(c)  $\frac{81}{256}$

(d)  $\frac{27}{1024}$

84. A coin is biased so that heads comes up thrice as likely as tails. In four independent tosses of the coin, what is probability of getting exactly three heads?

(a)  $\frac{81}{256}$

(b)  $\frac{27}{64}$

(c)  $\frac{27}{256}$

(d)  $\frac{9}{256}$

85. Let  $X$  and  $Y$  be two random variables such that  $X + Y = 100$ . If  $X$  follows Binomial distribution with parameters  $n = 100$  and  $p = \frac{4}{5}$ , what is the variance of  $Y$ ?

(a) 1

(b)  $\frac{1}{2}$

(c) 16

(d)  $\frac{1}{16}$

86. If two lines of regression are  $x + 4y + 1 = 0$  and  $4x + 9y + 7 = 0$ , then what is the value of  $x$  when  $y = -3$ ?

- (a) -13
- (b) -5
- (c) 5
- (d) 7

87. The central angles  $p$ ,  $q$ ,  $r$  and  $s$  (in degrees) of four sectors in a Pie Chart satisfy the relation  $9p = 3q = 2r = 6s$ . What is the value of  $4p - q$ ?

- (a) 12
- (b) 24
- (c) 30
- (d) 36

88. The observations 4, 1, 4, 3, 6, 2, 1, 3, 4, 5, 1, 6 are outputs of 12 dices thrown simultaneously. If  $m$  and  $M$  are means of lowest 8 observations and highest 4 observations respectively, then what is  $(2m + M)$  equal to?

- (a) 10
- (b) 12
- (c) 17
- (d) 21

89. A bivariate data set contains only two points  $(-1, 1)$  and  $(3, 2)$ . What will be the line of regression of  $y$  on  $x$ ?

- (a)  $x - 4y + 5 = 0$
- (b)  $3x + 2y - 1 = 0$

(c)  $x + 4y + 1 = 0$

(d)  $5x - 4y + 1 = 0$

90. A die is thrown 10 times and obtained the following outputs:

1, 2, 1, 1, 2, 1, 4, 6, 5, 4

What will be the mode of data so obtained?

- (a) 6
- (b) 4
- (c) 2
- (d) 1

Consider the following for the next three (03) items that follow:

Let  $I_1 = \int_0^\pi \frac{x}{1 + \cos^2 x} dx$  and

$I_2 = \int_0^\pi \frac{1}{1 + \sin^2 x} dx$

91. What is the value of  $\frac{I_1 + I_2}{I_1 - I_2}$ ?

- (a) 1
- (b)  $\pi$
- (c)  $\pi^2$
- (d)  $\frac{\pi + 1}{\pi - 1}$

92. What is the value of  $8I_1^2$ ?

- (a)  $\pi$
- (b)  $\pi^2$
- (c)  $\pi^3$
- (d)  $\pi^4$

93. What is the value of  $I_2$ ?

- (a)  $\frac{\pi}{\sqrt{2}}$
- (b)  $\frac{\pi}{2\sqrt{2}}$
- (c)  $\frac{3\pi}{2\sqrt{2}}$
- (d)  $\frac{\pi}{4\sqrt{2}}$

Consider the following for the next two (02) items that follow:

Let  $l = \int_a^b \frac{|x|}{x} dx$ ,  $a < b$

94. What is  $l$  equal to when  $a < 0 < b$ ?

- (a)  $a + b$
- (b)  $a - b$
- (c)  $b - a$
- (d)  $\frac{(a+b)}{2}$

95. What is  $l$  equal to when  $a < b < 0$ ?

- (a)  $a + b$
- (b)  $a - b$
- (c)  $b - a$
- (d)  $\frac{(a+b)}{2}$

Consider the following for the next three (03) items that follow:

Let  $f(x) = |\ln x|$ ,  $x \neq 1$

96. What is the derivative of  $f(x)$  at  $x = 0.5$ ?

- (a)  $-2$
- (b)  $-1$
- (c)  $1$
- (d)  $2$

97. What is the derivative of  $f(x)$  at  $x = 2$ ?

- (a)  $-\frac{1}{2}$
- (b)  $-1$
- (c)  $\frac{1}{2}$
- (d)  $2$



98. What is the derivative of  $f \circ f(x)$ , where  $1 < x < 2$  ?

- (a)  $\frac{1}{\ln x}$
- (b)  $\frac{1}{x \ln x}$
- (c)  $-\frac{1}{\ln x}$
- (d)  $-\frac{1}{x \ln x}$

Consider the following for the next two (02) items that follow :

$$\text{Let } f(x) = \begin{cases} x+6, & x \leq 1 \\ px+q, & 1 < x < 2 \\ 5x, & x \geq 2 \end{cases}$$

and  $f(x)$  is continuous

99. What is the value of  $p$  ?

- (a) 2
- (b) 3
- (c) 4
- (d) 5

100. What is the value of  $q$  ?

- (a) 2
- (b) 3
- (c) 4
- (d) 5

Consider the following for the next two (02) items that follow :

Consider the function

$$f(x) = |x-2| + |3-x| + |4-x|, \text{ where } x \in R.$$

101. At what value of  $x$  does the function attain minimum value ?

- (a) 2
- (b) 3
- (c) 4
- (d) 0

102. What is the minimum value of the function ?

- (a) 2
- (b) 3
- (c) 4
- (d) 0

Consider the following for the next two (02) items that follow :

Consider the sum

$$S = 0! + 1! + 2! + 3! + 4! + \dots + 100!$$

103. If the sum  $S$  is divided by 8, what is the remainder ?

- (a) 0
- (b) 1
- (c) 2
- (d) Cannot be determined

104. If the sum  $S$  is divided by 60, what is the remainder?

- (a) 1
- (b) 3
- (c) 17
- (d) 34

Consider the following for the next two (02) items that follow :

In a triangle  $PQR$ ,  $P$  is the largest angle and  $\cos P = \frac{1}{3}$ . Further the in-circle of the triangle touches the sides  $PQ$ ,  $QR$  and  $RP$  at  $N$ ,  $L$  and  $M$  respectively such that the lengths  $PN$ ,  $QL$  and  $RM$  are  $n$ ,  $n+2$ ,  $n+4$  respectively where  $n$  is an integer.

105. What is the value of  $n$ ?

- (a) 4
- (b) 6
- (c) 8
- (d) 10

106. What is the length of the smallest side?

- (a) 12
- (b) 14
- (c) 16
- (d) 18

Consider the following for the next two (02) items that follow :

Given that

$$\sin x + \cos x + \tan x + \cot x + \sec x + \operatorname{cosec} x = 7$$

107. The given equation can be reduced to

- (a)  $\sin^2 2x - 44 \sin 2x + 36 = 0$
- (b)  $\sin^2 2x + 44 \sin 2x - 36 = 0$
- (c)  $\sin^2 2x - 22 \sin 2x + 18 = 0$
- (d)  $\sin^2 2x + 22 \sin 2x - 18 = 0$

108. If  $\sin 2x = a - b\sqrt{c}$ , where  $a$  and  $b$  are natural numbers and  $c$  is prime number, then what is the value of  $a - b + 2c$ ?

- (a) 0
- (b) 14
- (c) 21
- (d) 28

Consider the following for the next two (02) items that follow :

A quadratic equation is given by

$$(3 + 2\sqrt{2})x^2 - (4 + 2\sqrt{3})x + (8 + 4\sqrt{3}) = 0$$

109. What is the HM of the roots of the equation?

- (a) 2
- (b) 4
- (c)  $2\sqrt{2}$
- (d)  $2\sqrt{3}$

110. What is the GM of the roots of the equation ?

(a)  $\sqrt{2}(\sqrt{6}-\sqrt{3}+\sqrt{2}-1)$

(b)  $\sqrt{2}(\sqrt{6}+\sqrt{3}-\sqrt{2}-1)$

(c)  $(\sqrt{6}-\sqrt{3}+\sqrt{2}-1)$

(d)  $(\sqrt{6}+\sqrt{3}+\sqrt{2}-1)$

Consider the following for the next two (02) items that follow :

Let  $\sin\beta$  be the GM of  $\sin\alpha$  and  $\cos\alpha$ ;  $\tan\gamma$  be the AM of  $\sin\alpha$  and  $\cos\alpha$ .

111. What is  $\cos 2\beta$  equal to ?

(a)  $(\cos\alpha - \sin\alpha)^2$

(b)  $(\cos\alpha + \sin\alpha)^2$

(c)  $(\cos\alpha - \sin\alpha)^3$

(d)  $\frac{(\cos\alpha - \sin\alpha)^2}{2}$

112. What is the value of  $\sec 2\gamma$ ?

(a)  $\frac{3 - \sin 2\alpha}{5 + 2 \sin 2\alpha}$

(b)  $\frac{5 + \sin 2\alpha}{3 - \sin 2\alpha}$

(c)  $\frac{3 - 2 \sin 2\alpha}{4 + \sin 2\alpha}$

(d)  $\frac{3 - \sin 2\alpha}{4 + 3 \sin 2\alpha}$

Consider the following for the next two (02) items that follow :

A flagstaff 20 m long standing on a pillar 10 m high subtends an angle  $\tan^{-1}(0.5)$  at a point  $P$  on the ground. Let  $\theta$  be the angle subtended by the pillar at this point  $P$ .

113. If  $x$  is the distance of  $P$  from bottom of the pillar, then consider the following statements :

1.  $x$  can take two values which are in the ratio 1 : 3

2.  $x$  can be equal to height of the flagstaff

Which of the statements given above is/are correct ?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

114. What is a possible value of  $\tan\theta$ ?

(a)  $\frac{3}{4}$

(b)  $\frac{2}{3}$

(c)  $\frac{1}{3}$

(d)  $\frac{1}{4}$

Consider the following for the next two (02) items that follow :

The perimeter of a triangle  $ABC$  is 6 times the AM of sine of angles of the triangle.

Further  $BC = \sqrt{3}$  and  $CA = 1$ .

115. What is the perimeter of the triangle ?

(a)  $\sqrt{3} + 1$

(b)  $\sqrt{3} + 2$

(c)  $\sqrt{3} + 3$

(d)  $2\sqrt{3} + 1$

116. Consider the following statements :

1.  $ABC$  is right angled triangle
2. The angles of the triangle are in AP

Which of the statements given above is/are correct ?

(a) 1 only

(b) 2 only

(c) Both 1 and 2

(d) Neither 1 nor 2

Consider the following for the next two (02) items that follow :

Let  $x = \frac{\sin^2 A + \sin A + 1}{\sin A}$  where  $0 < A \leq \frac{\pi}{2}$

117. What is the minimum value of  $x$  ?

(a) 1

(b) 2

(c) 3

(d) 4

118. At what value of  $A$  does  $x$  attain the minimum value ?

(a)  $\frac{\pi}{6}$

(b)  $\frac{\pi}{4}$

(c)  $\frac{\pi}{3}$

(d)  $\frac{\pi}{2}$

Consider the following for the next two (02) items that follow :

In the triangle  $ABC$ ,

$$a^2 + b^2 + c^2 = ac + \sqrt{3}bc$$

119. What is the nature of the triangle ?

(a) Equilateral

(b) Isosceles

(c) Right angled triangle

(d) Scalene but not right angled

120. If  $c = 8$ , what is the area of the triangle ?

(a)  $4\sqrt{3}$

(b)  $6\sqrt{3}$

(c)  $8\sqrt{3}$

(d)  $12\sqrt{3}$